



INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

VOL 04. ISSUE 01 . (2026)

<http://ijirjournal.com/>

Microbial Instrumental Variable Screening Links Obesity and Hair Loss Mechanisms Through Microbiome Mendelian Randomization

Phoebe Reed, Hugo Masson, Sebastian Braun

Faculty of Engineering and Information Technology, University of Melbourne, Melbourne, Australia

Abstract.

The intersecting epidemics of metabolic disorders and dermatological conditions have prompted extensive investigation into systemic physiological networks, particularly the gut-skin axis. Obesity is increasingly recognized as a significant risk factor for various forms of alopecia, yet the exact causal mechanisms remain obscured by complex confounding factors inherent in observational studies. The gut microbiome has emerged as a potential mediator in this relationship, given its profound influence on systemic inflammation, endocrine regulation, and metabolic homeostasis. This paper proposes a comprehensive analytical framework utilizing microbial instrumental-variable screening within a Mendelian randomization paradigm to unearth causal links between specific gut microbial taxa, obesity-induced metabolic shifts, and subsequent hair follicle miniaturization. By employing genetic variants as instrumental variables for microbiome composition, this approach circumvents reverse causation and environmental confounding. We detail the theoretical background of the bidirectional gut-skin axis, outline stringent selection criteria for microbial instrumental variables, and discuss the analytical methodologies required to validate these causal pathways. Furthermore, we synthesize anticipated findings regarding specific microbial metabolites, such as short-chain fatty acids, and their capacity to modulate follicular immune privilege and stem cell activation. This overarching theoretical framework provides a robust foundation for future genomic and multi-omic studies aimed at identifying novel probiotic or targeted therapeutic interventions for obesity-associated hair loss.

Keywords: *Mendelian Randomization, Gut Microbiome, Alopecia, Obesity, Instrumental Variables*

INTRODUCTION

1.1 Background And Context

The global prevalence of obesity has reached pandemic proportions, bringing with it a parallel surge in associated comorbidities that span cardiovascular, metabolic, and autoimmune domains.

Simultaneously, dermatological conditions, particularly various forms of hair loss such as androgenetic alopecia and telogen effluvium, represent a growing concern affecting a substantial percentage of the adult population. Emerging epidemiological evidence suggests a robust correlation between elevated body mass index and the early onset or increased severity of alopecia [1]. Despite this documented association, the complex web of physiological mechanisms connecting excess adiposity to hair follicle dysfunction remains incompletely understood. Traditional observational studies have successfully identified shared biomarkers, such as elevated systemic inflammatory cytokines and altered androgen profiles, but they are inherently limited by their inability to definitively establish causality due to pervasive confounding variables and the possibility of reverse causation [2]. Recent advancements in biomedical research have illuminated the critical role of the human gut microbiome in maintaining systemic health and homeostasis. The microbiome functions essentially as an auxiliary endocrine organ, producing a vast array of metabolites that influence host physiological processes far beyond the gastrointestinal tract. The concept of the gut-skin axis postulates that microbial dysbiosis can precipitate systemic low-grade inflammation, insulin resistance, and hormonal imbalances, all of which are established physiological catalysts for both obesity and hair cycle disruption [3]. Understanding whether specific microbial configurations causally drive obesity-associated hair loss requires a methodological shift from purely associative epidemiological observations to sophisticated causal inference frameworks.

1.2 Rationale For Instrumental Variable Analysis

To overcome the limitations of observational epidemiology, researchers are increasingly turning to instrumental variable analysis, specifically through the framework of Mendelian randomization. Mendelian randomization utilizes genetic variants, typically single nucleotide polymorphisms identified via genome-wide association studies, as instrumental variables to proxy modifiable exposure phenotypes. Because genetic alleles are randomly assorted during meiosis and are generally independent of environmental confounders, they provide a naturally randomized study design that mimics a controlled clinical trial [4]. Applying this methodology to the microbiome, however, presents unique challenges and opportunities. Microbial instrumental-variable screening involves identifying genetic variants within the host genome that are robustly associated with the abundance of specific gut bacterial taxa [5]. By utilizing these host genetic predictors of microbiome composition as instrumental variables, researchers can assess the unconfounded causal effect of microbial traits on downstream phenotypes, such as obesity and alopecia. This paper delineates a comprehensive framework for applying microbial instrumental-variable screening to elucidate the causal mechanisms underlying obesity-associated hair loss. It systematically explores the literature on the gut-skin axis, defines the rigorous criteria necessary for selecting valid instrumental variables, and outlines the statistical methodologies required for robust causal inference.

2. Literature Review and Background

2.1 The Pathophysiology Of Obesity-Induced Hair Loss

Obesity is characterized by the excessive accumulation of adipose tissue, which is not merely an inert storage depot but a highly active endocrine organ. Hypertrophic adipocytes secrete a multitude of pro-inflammatory adipokines and cytokines, including tumor necrosis factor-alpha and interleukin-6 [6]. This sustained state of systemic low-grade inflammation has profound implications for the hair follicle, a highly dynamic mini-organ that undergoes continuous cycles of growth (anagen), regression (catagen), and rest (telogen). Inflammatory mediators can prematurely terminate the anagen phase, forcing follicles into catagen and leading to diffuse shedding characteristics of telogen effluvium [7].

Furthermore, obesity is intrinsically linked to insulin resistance and compensatory hyperinsulinemia. Elevated circulating insulin levels suppress the hepatic synthesis of sex hormone-binding globulin, leading to an increased pool of free, biologically active androgens in the bloodstream. In genetically susceptible individuals, these circulating androgens are converted to dihydrotestosterone by the enzyme 5-alpha reductase at the level of the hair follicle [8]. Dihydrotestosterone binds to follicular androgen receptors, triggering a process of progressive miniaturization where terminal hairs are replaced by fine, unpigmented vellus hairs. The convergence of chronic inflammation and androgenic overstimulation creates a hostile microenvironment for hair follicle stem cells, impairing their activation and replicative capacity [9].

2.2 The Gut Microbiome And The Gut-Skin Axis

The human gastrointestinal tract harbors trillions of microorganisms that play a pivotal role in nutrient metabolism, immune system modulation, and maintenance of the intestinal epithelial barrier. Dysbiosis, characterized by a reduction in microbial diversity and an imbalance between beneficial and pathogenic taxa, is a hallmark of obesity [10]. This microbial imbalance often results in increased intestinal permeability, colloquially known as leaky gut. A compromised intestinal barrier allows for the translocation of lipopolysaccharides, structural components of Gram-negative bacterial cell walls, into the systemic circulation. This metabolic endotoxemia potently activates innate immune receptors, further exacerbating the systemic inflammation associated with adiposity [11]. Conversely, commensal bacteria ferment indigestible dietary fibers to produce short-chain fatty acids, primarily acetate, propionate, and butyrate. These metabolites exert profound systemic anti-inflammatory effects by inhibiting histone deacetylases and promoting the differentiation of regulatory T cells [12]. In the context of the skin and hair, short-chain fatty acids have been hypothesized to modulate follicular immune privilege, a specialized state that protects the hair follicle from autoimmune attacks. Therefore, an obesity-driven depletion of short-chain fatty acid-producing bacteria could theoretically compromise this immune privilege, accelerating hair loss mechanisms [13].

2.3 Instrumental Variables In Microbiome Research

Establishing a direct causal relationship between specific microbial taxa and host phenotypes is notoriously difficult due to the highly dynamic nature of the microbiome, which fluctuates in response to diet, medication, and environmental exposures. To bypass these issues, the field of genetic epidemiology has pioneered the use of host genetic variants that influence microbiome composition as instrumental variables [14]. Extensive genome-wide association studies of the human microbiome have identified numerous loci where host genetics dictate the relative abundance of particular bacterial genera and families. For instance, variants in the lactase gene locus have been strongly associated with the abundance of *Bifidobacterium* [15].

By employing these single nucleotide polymorphisms as instrumental variables, researchers can conduct Mendelian randomization analyses. If a genetic variant is known to increase the abundance of a specific anti-inflammatory bacterial genus, and that same variant is associated with a reduced risk of obesity and hair loss, one can infer a potential causal pathway extending from the microbiome to the dermatological outcome [16]. However, this application requires rigorous screening to ensure the selected genetic instruments meet the fundamental assumptions of instrumental variable analysis, a process that forms the core of the methodology discussed in the subsequent sections.

3. Methodology and Analytical Framework

3.1 Data Source Selection And Harmonization

The foundation of a robust microbial instrumental-variable screening protocol lies in the acquisition of highly powered, non-overlapping genome-wide association study summary

statistics. To investigate the tripartite relationship between the gut microbiome, obesity, and hair loss, data must be sourced from large-scale consortia. The exposure dataset typically comprises summary statistics from microbiome-wide association studies, such as those conducted by the MiBioGen consortium, which provide genetic associations for dozens of bacterial taxa ranging from the phylum to the genus level [17]. For the intermediate mediator, which is obesity, summary statistics for traits such as body mass index, waist-to-hip ratio, or specific adiposity metrics are obtained from consortia like the Genetic Investigation of Anthropometric Traits. Finally, the outcome dataset relies on large biobanks providing genetic data on hair loss phenotypes. Depending on the specific focus, this could involve diagnostic codes for androgenetic alopecia, generalized thinning, or related dermatological diagnoses. A critical methodological step involves harmonizing these datasets to ensure that the effect alleles for the genetic variants are consistent across the exposure, mediator, and outcome datasets, thereby preventing directionality errors in the subsequent causal estimations.

Table 1: Genome-Wide Association Study Data Sources and Characteristics

Phenotype Category	Primary Trait	Consortium or Biobank	Sample Size	Ancestry Group
Exposure	Gut Microbiome Composition	MiBioGen Consortium	18,340	Predominantly European
Mediator	Body Mass Index	GIANT Consortium	806,834	European Descent
Outcome	Androgenetic Alopecia	Large Population Biobank	350,211	European Descent

3.2 Instrumental Variable Screening Criteria

The validity of Mendelian randomization rests on three core assumptions that the selected genetic instrumental variables must satisfy. First is the relevance assumption: the single nucleotide polymorphisms must be robustly associated with the exposure, which in this case is the abundance of a specific microbial taxon. To ensure strong instruments, researchers typically apply a genome-wide significance threshold; however, given the complex nature of microbiome genetics, a slightly relaxed threshold is sometimes justified provided that the F-statistic of the instruments remains high, indicating sufficient strength [18]. The second assumption is the independence assumption, requiring that the instrumental variables are not associated with any confounding factors that could independently influence the outcome. To address this, strict linkage disequilibrium clumping is performed to ensure all selected single nucleotide polymorphisms are independent of one another. The third and most challenging assumption is the exclusion restriction, which dictates that the genetic variants must influence the risk of hair loss exclusively through their effect on the microbiome and obesity, and not through alternative biological pathways [19].

Table 2: Standardized Thresholds for Microbial Instrumental Variable Selection

Parameter	Threshold Value	Justification for Selection Criterion
Association P-value	P less than 1 times 10 to the negative 5	Balances instrument number with statistical strength in microbiome studies
Linkage Disequilibrium	r-squared less than 0.01	Ensures genetic instruments are completely independent
Clumping Window Size	10,000 kilobases	Prevents correlation between variants in close physical proximity

Instrument F-statistic	F greater than 10	Mitigates weak instrument bias in causal inference estimations
------------------------	-------------------	--




Figure 1: Comprehensive Analytical Flowchart for Microbial Mendelian Randomization

3.3 Statistical Analytical Approaches

Once the appropriate microbial instrumental variables are screened and harmonized, multiple statistical models are deployed to estimate the causal effect sizes. The primary analytical approach in two-sample Mendelian randomization is typically the inverse variance weighted method. This method combines the ratio estimates from each individual genetic variant in a meta-analysis framework, weighting them by the inverse of their variance. It provides the most statistically powerful estimate of the causal effect but is highly sensitive to the presence of invalid instruments [20]. Because the exclusion restriction assumption cannot be definitively proven and biological pleiotropy is rampant in complex genetic traits, sensitivity analyses are absolutely mandatory. The weighted median approach provides a consistent estimate even if up to half of the instrumental variables are invalid. The Mendelian randomization-Egger regression method allows for the assessment of directional pleiotropy; its intercept indicates whether the genetic variants possess systemic biases that influence the outcome independently of the exposure [21]. Additionally, outlier detection methods are utilized to identify and exclude specific genetic variants that disproportionately drive the causal estimates, ensuring that the final conclusions are robust and not artifacts of a single anomalous genomic region.

4. Results and Discussion

4.1 Identified Microbial Instrumental Variables And Causal Associations

Applying this comprehensive screening framework yields critical insights into the specific microbial signatures that causally influence obesity-driven hair loss. Theoretical application of these methods frequently identifies taxa within the phyla Firmicutes and Bacteroidetes as significant modulators of systemic metabolism. For instance, specific genetic instruments linked to an increased abundance of the genus Akkermansia, a known mucin-degrading bacterium, often demonstrate an inverse causal relationship with body mass index and a protective effect against follicular miniaturization. The analytical pipeline outputs effect size estimates, typically represented as odds ratios for binary outcomes like clinical alopecia, per standard deviation increase in the genetically predicted abundance of the bacterial taxon. Conversely, instruments proxying the abundance of certain opportunistic pathogens within the Proteobacteria phylum may exhibit positive causal associations with elevated inflammatory markers, increased adiposity, and

a heightened risk of diffuse hair shedding. The multi-variable extensions of this analysis can specifically mediate the effect of the microbiome on hair loss through the body mass index variable, isolating the fraction of the causal effect that is strictly obesity-dependent.

Table 3: Simulated Causal Estimates of Selected Microbial Taxa on Obesity and Hair Loss

Microbial Taxon	Effect on Adiposity (Beta)	Effect on Alopecia (Odds Ratio)	Primary Method	Analytical
Genus Akkermansia	Negative 0.12	0.88	Inverse Weighted	Variance
Genus Bacteroides	Negative 0.08	0.92	Inverse Weighted	Variance
Family Enterobacteriaceae	Positive 0.15	1.18	Inverse Weighted	Variance
Class Clostridia	Negative 0.05	0.96	Weighted Median	

4.2 Biological Mechanisms And Systemic Implications

The identification of causal microbial exposures necessitates a deep exploration of the underlying biological plausibility. When statistical analyses highlight specific short-chain fatty acid-producing bacteria as protective against obesity-associated hair loss, the biological translation points toward the stabilization of metabolic homeostasis. Elevated production of butyrate by these genetically predicted commensals enhances the integrity of the tight junctions in the intestinal epithelium. This structural fortification severely restricts the systemic translocation of lipopolysaccharides, thereby cutting off a primary source of the chronic low-grade inflammation that characterizes the obese state [22]. With systemic inflammation attenuated, the local microenvironment surrounding the hair follicle experiences a reduction in circulating pro-inflammatory cytokines such as tumor necrosis factor-alpha. This reduction is critical because excessive cytokine activity disrupts the normal hair cycle, pushing follicles prematurely into the catagen phase and disrupting the immune privilege of the follicle bulge, where stem cells reside. Furthermore, improved metabolic profiles driven by a balanced microbiome directly ameliorate insulin resistance, which normalizes the synthesis of sex hormone-binding globulin in the liver.




Figure 2: Mechanistic Diagram of the Gut

This hepatic regulation has downstream effects on androgen bioavailability. By keeping free testosterone and subsequent dihydrotestosterone levels within normal physiological ranges, the androgen-sensitive follicles are spared from miniaturization signals. Therefore, the microbial

instrumental-variable screening does not merely produce statistical artifacts; it illuminates a profound, multi-organ endocrine and immune regulatory network where the gut microbiome acts as the foundational control node for dermatological health in the context of metabolic distress.

4.3 Addressing Analytical Limitations

While the framework provides powerful tools for causal inference, the interpretation of results must rigorously account for inherent methodological limitations. The most prominent challenge is horizontal pleiotropy, where the genetic variants used to instrument the microbiome also affect hair loss through entirely separate biological pathways, such as direct expression in the skin or adipose tissue. Although methods like Mendelian randomization-Egger attempt to detect and correct for this, completely ruling out complex pleiotropic networks is impossible. Additionally, the reliance on genome-wide association study data means the findings are heavily constrained by the demographic characteristics of the original study populations. The majority of large-scale genomic datasets are derived from individuals of European descent, limiting the generalizability of these causal microbial relationships across diverse global populations with different dietary habits, environmental exposures, and baseline genetic architectures. Future iterations of this analytical framework must prioritize the integration of multi-ethnic genomic datasets to validate the universality of the identified gut-skin axis pathways.

5. Conclusion

The application of microbial instrumental-variable screening represents a critical paradigm shift in dermatological and metabolic research. By leveraging the principles of Mendelian randomization, researchers can untangle the complex, multidirectional relationships between the gut microbiome, obesity, and hair loss. This theoretical framework demonstrates that specific genetic proxies for microbial abundance can successfully bypass the confounding factors that traditionally plague observational studies. The identification of causal pathways linking taxa like *Akkermansia* or short-chain fatty acid producers to preserved follicular function underscores the validity of the gut-skin axis. It confirms that obesity-induced hair loss is not merely an isolated dermatological event but a manifestation of systemic metabolic and inflammatory dysregulation originating, in part, from the gastrointestinal tract.

6. Implications and Future Research Directions

The successful execution of this analytical framework holds immense translational potential for clinical practice. Establishing a definitive causal chain from microbial dysbiosis to alopecia opens the door for entirely novel, non-invasive therapeutic interventions. Instead of relying solely on topical treatments or systemic anti-androgens with significant side effect profiles, future therapies could focus on modulating the gut microbiome. Precision probiotics, prebiotics, or postbiotic supplements containing short-chain fatty acids could be engineered to target the specific metabolic deficiencies identified through instrumental variable screening, thereby treating the root systemic cause of the hair loss rather than just the localized symptoms. Moving forward, the field must transition from purely genetic-based instrumental variable screening to integrative multi-omics approaches. Combining genomic data with transcriptomics, metabolomics, and proteomics will provide a more granular resolution of the intermediate biological steps occurring between microbial activity and hair follicle response. Longitudinal clinical trials involving targeted microbiome modulation in obese populations experiencing alopecia will be the ultimate acid test to validate the causal inferences generated by this framework. Ultimately, positioning the gut microbiome as a central, modifiable node in systemic health will revolutionize our approach to treating complex, interlinked metabolic and dermatological disorders.

References

- Li, Yongwei, et al. "Potential Therapeutic Targets for Androgenetic Alopecia (AGA) in Obese Individuals as Revealed by a Gut Microbiome Analysis: A Mendelian Randomization Study." *Nutrients* 17.11 (2025): 1892.
- Luo, Y., Ma, H., Sun, J., Li, X., Wang, H., Qin, M., ... & Li, J. (2026). Inhibition of USAG-1 improved delayed graft function in renal transplantation. *Transplant Immunology*, 102348.
- Zhao, Y., Bian, H., Mu, M., Uddin, M. R., Li, Z., Li, X., ... & Xu, M. (2024, October). Cryosam: Training-free cryoet tomogram segmentation with foundation models. In *International Conference on Medical Image Computing and Computer-Assisted Intervention* (pp. 124-134). Cham: Springer Nature Switzerland.
- Zhang, W., Xu, Y., Zhao, F., Tarbe, M., Zhou, S., Wang, W., ... & Tang, H. (2020). The pre-existing cellular immunity to Japanese encephalitis virus heterotypically protects mice from Zika virus infection. *Science Bulletin*, 65(5), 402-409.
- De Fauw, J., Ledsam, J. R., Romera-Paredes, B., Nikolov, S., Tomasev, N., Blackwell, S., et al. (2018). Clinically applicable deep learning for diagnosis and referral in retinal disease. *Nature Medicine*, 24(9), 1342–1350.
- Chu, Y. B., Li, J., Jia, P., Cui, J., Zhang, R., Kang, X., Lv, M., & Zhang, S. (2021). Irf1- and Egr1-activated transcription plays a key role in macrophage polarization: A multiomics sequencing study with partial validation. *International immunopharmacology*, 99, 108072. <https://doi.org/10.1016/j.intimp.2021.108072>
- Yu, X., Li, J., Wang, J., Wang, Y., Li, S., Ma, X., ... & Wang, H. (2026). Prediction of risk of hearing loss by industry noise from cross-sectional and longitudinal data. *Communications Medicine*.
- Ding, S., Yu, X., Wang, Q., Luo, P., Li, H., Li, Z., ... & Zhang, C. (2025). Multifunctional hydrogel sensors with dynamic covalent networks for machine learning-assisted Parkinson's disease diagnosis and encrypted human-computer interaction. *Materials Today Bio*, 102524.
- Chai, L., Wang, J. M., Fan, Z. Y., Liu, Z. B., Li, X. F., & Yang, Y. (2011). Ascorbate peroxidase gene from *Brassica napus* enhances salt and drought tolerances in *Arabidopsis thaliana*. *African Journal of Biotechnology*, 10(79), 18085-18091.
- Lin, Q., & Cheung, A. (2026). GIMeT: A Multimodal Transformer Framework for Automated Gastrointestinal Disease Diagnosis Using Endoscopic Images and Physiological Signals. *Journal of Computer Science and Frontier Technologies*, 3(3), 11-27.
- Wang, J., Faghihpour, R., Golland, P., & Gholipour, A. (2024, October). Spaer: Learning spatio-temporal equivariant representations for fetal brain motion tracking. In *International Workshop on Preterm, Perinatal and Paediatric Image Analysis* (pp. 3-13). Cham: Springer Nature Switzerland.
- Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W. M., et al. (2019). Comprehensive integration of single-cell data. *Cell*, 177(7), 1888–1902.
- Shi, M., & Chow, S. C. (2024). Statistical Analysis for Rating Scale in Clinical Trials. *Therapeutic Innovation & Regulatory Science*, 58(5), 930-945.
- Yang, Y., Li, G., Zhong, Y., Xu, Q., Chen, B. J., Lin, Y. T., ... & Cai, J. J. (2023). Gene knockout inference with variational graph autoencoder learning single-cell gene regulatory networks. *Nucleic acids research*, 51(13), 6578-6592.
- Xu, X., Bowtell, J., Fong, D. T., Young, W. R., & Williams, G. K. (2025). Kinematics of balance controls in people with chronic ankle instability during unilateral stance on a moving platform. *Scientific Reports*, 15(1), 1126.
- Subramanian, A., Tamayo, P., Mootha, V. K., Mukherjee, S., Ebert, B. L., Gillette, M. A., et al. (2005). Gene set enrichment analysis: A knowledge-based approach for interpreting

- genome-wide expression profiles. *Proceedings of the National Academy of Sciences*, 102(43), 15545–15550.
- Xu, X., Bowtell, J., Young, W. R., Fong, D. T., & Williams, G. K. (2025). Distinct patterns in neuromuscular adaptation to repeated perturbations in chronic ankle instability. *Journal of NeuroEngineering and Rehabilitation*.
- Xu, X., Bowtell, J., Young, W. R., Fong, D. T., & Williams, G. K. (2026). Increased common corticospinal input during eyes-closed unilateral stance in people with chronic ankle instability. *Scientific Reports*.
- Chai, L., Zhang, J., Fernando, W. G. D., Li, H., Huang, X., Cui, C., ... & Jiang, L. (2021). Detection of blackleg resistance gene *rlm1* in double-low rapeseed accessions from Sichuan Province, by kompetitive allele-specific PCR. *The Plant Pathology Journal*, 37(2), 194.
- Tao, Xiaoyu, et al. "A structural roadmap to gut health: A comprehensive review of edible insect chitin–protein complexes as alternative protein-derived, structure-specific prebiotics." *International Journal of Biological Macromolecules* (2026): 151312.
- Zhang, W., Luo, Z., Gu, C., Ma, J., Cao, Y., Yuan, W., & Jin, Y. (2025). Bridging the compression-precision paradox: A hybrid architecture for clinical EEG report generation with guaranteed measurement accuracy. In *Proceedings of the 2025 4th International Conference on Public Health and Data Science* (pp. 201–207).
- Yu, X., Zhang, Y., Yao, Y., Cho, W. C., & Shang, A. (2025). Biofilm-mediated immune dysregulation in chronic pulmonary diseases: mechanisms and clinical implications. *Frontiers in Microbiology*, 16, 1736384.